



Kansas City

An Independent Licensee of the Blue Cross and Blue Shield Association

In Vitro Chemoresistance and Chemosensitivity Assays

Policy Number: 2.03.01

Last Review: 10/2018

Origination: 10/1988

Next Review: 10/2019

Policy

Blue Cross and Blue Shield of Kansas City (Blue KC) will not provide coverage for In Vitro Chemoresistance and Chemosensitivity Assays. This is considered investigational.

When Policy Topic is covered

Not Applicable.

When Policy Topic is not covered

In vitro chemosensitivity assays, including but not limited to the Histoculture Drug Response Assay, a fluorescent cytoprint assay, the ChemoFx assay, are considered **investigational**.

In vitro chemoresistance assays, including but not limited to extreme drug resistance assays, are considered **investigational**.

Description of Procedure or Service

Populations	Interventions	Comparators	Outcomes
Individuals: • With cancer who are initiating chemotherapy	Interventions of interest are: • Chemosensitivity assays	Comparators of interest are: • Chemotherapy selection without chemosensitivity assay	Relevant outcomes include: • Overall survival • Disease-specific survival • Test accuracy • Test validity • Quality of life
Individuals: • With cancer who are initiating chemotherapy	Interventions of interest are: • Chemoresistance assays	Comparators of interest are: • Chemotherapy selection without chemoresistance assay	Relevant outcomes include: • Overall survival • Disease-specific survival • Test accuracy • Test validity • Quality of life

In vitro chemoresistance and chemosensitivity assays have been developed to provide information about the characteristics of an individual patient’s malignancy to predict potential responsiveness of their cancer to specific drugs. Thus, these assays are sometimes used by oncologists to select treatment regimens for an

individual patient. Several assays have been developed that differ with respect to processing of biological samples and detection methods. However, all involve similar principles and they share protocol components including: 1) isolation of cells and establishment in an in vitro medium (sometimes in soft agar); 2) incubation of the cells with various drugs; 3) assessment of cell survival; and 4) interpretation of the result.

For individuals who have cancer who are initiating chemotherapy who receive chemosensitivity assays, the evidence includes correlational observational studies. Relevant outcomes are overall survival, disease-specific survival, test accuracy and validity, and quality of life. Some retrospective and prospective correlational studies have suggested that chemoresistance assays may be associated with chemotherapy response. However, prospective studies do not consistently demonstrate that chemoresistance assay results are associated with survival. Furthermore, no studies were identified that compared outcomes for patients managed with assay-directed therapy to those managed with physician-directed therapy. Large, randomized, prospective clinical studies comparing clinical outcomes are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have cancer who are initiating chemotherapy who receive chemosensitivity assays, the evidence includes 1 randomized controlled trial (RCT) and correlational observational studies. Relevant outcomes are overall survival, disease-specific survival, test accuracy and validity, and quality of life. The most direct evidence on the effectiveness of chemosensitivity assays in the management of patients with cancer comes from several studies comparing outcomes for patients managed with a chemosensitivity assay to those managed with standard care, including 1 RCT. Although some improvements in tumor response were noted, there were no differences in survival outcomes. A number of retrospective and prospective studies of several different chemosensitivity assays have suggested that patients whose tumors have higher chemosensitivity have better outcomes. Currently, additional studies to determine whether the clinical use of in vitro chemosensitivity testing leads to better outcomes are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

Background

A variety of chemosensitivity and chemoresistance assays have been clinically evaluated in human trials. All assays use characteristics of cell physiology to distinguish between viable and non-viable cells to quantify cell kill following exposure to a drug of interest. With few exceptions, drug doses used in the assays are highly variable depending on tumor type and drug class, but all assays require drug exposures ranging from several-fold below physiological relevance to several-fold above physiological relevance. Although a variety of assays exist to examine chemosensitivity or chemoresistance, only a few are commercially available. Available assays are outlined as follows:

Methods using differential staining/dye exclusion:

- The Differential Staining Cytotoxicity assay.(1) This assay relies on dye exclusion of live cells after mechanical disaggregation of cells from surgical or biopsy specimens by centrifugation. Cells are then established in culture and treated with the drugs of interest at 3 dose levels; the middle dose is that which could be achieved in therapy; 10-fold lower than the physiologically relevant dose; and, 10-fold higher. Exposure time ranges from 4 to 6 days; then, cells are restained with fast green dye and counterstained with hematoxylin and eosin (H&E). The fast green dye is taken up by dead cells, and H&E can then differentiate tumor cells from normal cells. The intact cell membrane of a live cell precludes staining with the green dye. Drug sensitivity is measured by the ratio of live cells in the treated samples to the number of live cells in the untreated controls.
- The EVA/PCD™ assay (available from Rational Therapeutics). This assay relies on ex vivo analysis of programmed cell death, as measured by differential staining of cells after apoptotic and nonapoptotic cell death markers in tumor samples exposed to chemotherapeutic agents. Tumor specimens obtained through biopsy or surgical resection are disaggregated using DNase and collagenase IV to yield tumor clusters of the desired size (50-100 cell spheroids). Because these cells are not proliferated, these microaggregates are believed to more closely approximate the human tumor microenvironment. These cellular aggregates are treated with the dilutions of the chemotherapeutic drugs of interest and incubated for 3 days. After drug exposure is completed, a mixture of Nigrosin B & Fast green dye with glutaraldehyde-fixed avian erythrocytes is added to the cellular suspensions.(2) The samples are then agitated and cytospin-centrifuged and, after air drying, are counterstained with H&E. The end point of interest for this assay is cell death, as assessed by observing the number of cells differentially stained due to changes in cellular membrane integrity.(3)
- The fluorometric microculture cytotoxicity assay is another cell viability assay that relies on the measurement of fluorescence generated from cellular hydrolysis of fluorescein diacetate to fluorescein in viable cells.(4) Cells from tumor specimens are incubated with cytotoxic drugs; drug resistance is associated with higher levels of fluorescence.

Methods using incorporation of radioactive precursors by macromolecules in viable cells:

- Tritiated thymine incorporation measures uptake of tritiated thymidine by DNA of viable cells. Using proteases and DNase to disaggregate the tissue, samples are seeded into single-cell suspension cultures on soft agar. They are then treated with the drug(s) of interest for 4 days. After 3 days, tritiated thymidine is added. After 24 hours of additional incubation, cells are lysed, and radioactivity is quantified and compared with a blank control consisting of cells that were treated with sodium azide. Only cells that are viable and proliferating will take up the radioactive thymidine. Therefore, there is an inverse relationship between uptake of radioactivity and sensitivity of the cells to the agent(s) of interest.(5)

- The Oncotech Extreme Drug Resistance EDR® assay (Exiqon Diagnostics; no longer commercially available) is methodologically similar to the thymidine incorporation assay, using metabolic incorporation of tritiated thymidine to measure cell viability; however, single cell suspensions are not required, so the assay is simpler to perform.⁶ Tritiated thymidine is added to the cultures of tumor cells, and uptake is quantified after various incubation times. Only live (resistant) cells will incorporate the compound. Therefore, the level of tritiated thymidine incorporation is directly related to chemoresistance. The interpretation of the results is unique in that resistance to the drugs is evaluated, as opposed to evaluation of responsiveness. Tumors are considered to be highly resistant when thymidine incorporation is at least 1 standard deviation above reference samples.

Methods to quantify cell viability by colorimetric assay:

- The Histoculture Drug Resistance Assay (HDRA; AntiCancer Inc., San Diego, CA).⁽⁷⁾ This assay evaluates cell growth after chemotherapy treatment based on a colorimetric assay that relies on mitochondrial dehydrogenases in living cells. Drug sensitivity is evaluated by quantification of cell growth in the 3-dimensional collagen matrix. There is an inverse relationship between the drug sensitivity of the tumor and cell growth. Concentrations of drug and incubation times are not standardized and vary depending on drug combination and tumor type.

Methods using incorporation of chemoluminescent precursors by macromolecules in viable cells:

- The Adenosine Triphosphate (ATP) Bioluminescence assay. This assay relies on measurement of ATP to quantify the number of viable cells in a culture. Single cells or small aggregates are cultured, and then exposed to drugs. Following incubation with drug, the cells are lysed and the cytoplasmic components are solubilized under conditions that will not allow enzymatic metabolism of ATP. Luciferin and firefly luciferase are added to the cell lysis product. This catalyzes the conversion of ATP to adenosine di- and monophosphate, and light is emitted proportionally to metabolic activity. This is quantified with a luminometer. From the measurement of light, the number of cells can be calculated. A decrease in ATP indicates drug sensitivity, whereas no loss of ATP suggests that the tumor is resistant to the agent of interest.
- The ChemoFX (Helomics, previously called Precision Therapeutics) assay also relies on quantifying ATP-based on chemoluminescence.^(8,9) Cells must be grown in a monolayer rather than in a 3-dimensional matrix.

Regulatory Status

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments. Chemoresistance and chemosensitivity assays discussed in this review are available under the auspices of the Clinical Laboratory Improvement Amendments. Laboratories that offer laboratory-developed tests must be licensed by the Clinical Laboratory Improvement Amendments for high-complexity testing. To date, the U.S. Food

and Drug Administration has chosen not to require any regulatory review of this test.

Rationale

This evidence review was created in March 1996 and has been updated regularly with searches of the MEDLINE database. The most recent literature update was performed through May 7, 2018.

This review was informed by a TEC Assessment (2002)¹⁰ and a systematic review by Samson et al (2004),¹¹ which concluded collectively that the evidence was insufficient to support the use of chemoresistance and chemosensitivity assays for guiding the choice of therapy regimen in cancer patients.

A variety of studies have reported a correlation between in vitro prediction or response and clinical response. While these studies may have internal validity, they cannot answer whether patients given assay-guided therapy or empirical therapy have different outcomes. To determine whether assay-guided treatment results in overall different outcomes than empirical treatment, it is important to take into account response rates, survival, adverse events, and quality of life (QOL). These effects may be assessed indirectly (eg, using decision analysis) or directly with comparative trials. Both the 2002 TEC Assessment and the 2004 systematic review^{10,11} recommended validating chemotherapy resistance and sensitivity assays with direct evidence gathered from prospective trials comparing patients treated empirically with patients treated with assay-directed therapy. In this way, not only can response rates and survival be taken into account, but also adverse events (eg, from the toxic effects of an ineffective drug or delay or loss of benefits of an effective drug) and QOL.

Evidence reviews assess whether a medical test is clinically useful. A useful test provides information to make a clinical management decision that improves the net health outcome. That is, the balance of benefits and harms is better when the test is used to manage the condition than when another test or no test is used to manage the condition.

The first step in assessing a medical test is to formulate the clinical context and purpose of the test. The test must be technically reliable, clinically valid, and clinically useful for that purpose. Evidence reviews assess the evidence on whether a test is clinically valid and clinically useful. Technical reliability is outside the scope of these reviews, and credible information on technical reliability is available from other sources.

Chemoresistance and chemosensitivity Assays

Clinical Context and Test Purpose

The purpose of chemoresistance and chemosensitivity assays in patients who have cancer is to inform the decision on selecting treatment regimens.

The question addressed in this evidence review is: Does the use of chemoresistance and chemosensitivity assays improve the net health outcome in individuals with cancer?

The following PICOTS were used to select literature to inform this review.

Patients

The relevant population of interest is individuals with cancer.

Interventions

The tests being considered are chemoresistance and chemosensitivity assays. These assays differ with respect to the processing of biologic samples and detection methods, but all involve (1) isolation of cancer cells and establishment in an in vivo medium; (2) incubation of cells with various drugs; (3) assessment of cell survival; and (4) interpretation of results.

Comparators

The following practice is currently being used: physician-choice of chemotherapy, without information from chemoresistance assays or chemosensitivity assays.

Outcomes

The general outcome of interest is to determine cancer cell viability following exposure to a cancer drug. An accurate determination of cancer cell viability to a particular drug can then inform the decision on which drug treatment will be most effective. The specific outcome of interest is survival (overall and/or progression-free). Additional relevant clinical outcomes are avoidance of chemotherapy-related toxicity and improved QOL.

False-positive and false-negative test results can lead to the patient receiving a treatment regimen that is potentially not the most effective.

Timing

Chemoresistance and chemosensitivity assays can be performed on fresh tumor tissue samples of newly diagnosed or recurrent cancer patients.

Setting

Cancer cells are obtained through clinically indicated surgery or biopsy. The samples are shipped to a laboratory conducting the chemoresistance or chemosensitivity assays.

Chemoresistance Assays

Chemoresistance assays are used to deselect potential chemotherapeutic regimens. The negative predictive value (NPV) is a key statistical measure. Unless the NPV is high, there is a chance that clinical decision making based on a chemoresistance assay could inappropriately exclude an effective therapy. The NPV will vary according to the prior probability of chemoresistance as well as the assay's sensitivity and specificity. The TEC Assessment (2002) concluded that chemoresistance assays have the highest clinical relevance in tumors with a low

probability of response.¹⁰ The Extreme Drug Resistance (EDR) assay was specifically designed to produce a very high NPV (>99%), such that the possibility of inappropriately excluding effective chemotherapy is remote in all clinical situations.¹²

The bulk of the literature on EDR assays has focused on correlational studies assessing results from predictive in vitro assays and observed outcomes of chemotherapy. However, in these studies, patients did not receive assay-guided chemotherapy regimens. As discussed in the systematic review by Samson et al (2004), correlational studies are inadequate to demonstrate the clinical utility of chemoresistance assays for several reasons.¹¹ First, such studies often aggregate patients with different tumor types, disease characteristics, chemotherapy options, and probabilities of response. This process is problematic because the accuracy of each assay used to predict in vivo response probably varies across different malignancies and patient characteristics. Second, the method by which assay results are translated into treatment decisions is not standardized. Third, it is important to consider not only response but also survival, QOL, and adverse events. The overall value of assay-guided therapy depends on the net balance of all health outcomes observed after treatment for all patients subjected to testing, regardless of assay results or the accuracy of its predicted response. Examples of some of the earlier published correlation studies of the EDR assay include those by Eltabbakh et al (1998, 2000),^{13,14} Mehta et al (2001),¹⁵ Holloway et al (2002),¹⁶ and Ellis et al (2002).¹⁷

The TEC Assessment (2002) identified a nonrandomized retrospective comparative study by Loizzi et al (2003)¹⁸ using the EDR assay. (While this study of patients with recurrent ovarian cancer found a significantly higher overall response rate, better progression-free survival (PFS), and higher overall survival (OS) among platinum-sensitive patients receiving assay-guided therapy, it was not designed to address potential biases and confounding adequately. Since the Loizzi study, no additional comparative studies of assay-guided therapy vs physician-directed therapy have appeared for chemoresistance assays.

Technically Reliable

Assessment of technical reliability focuses on specific tests and operators and requires review of unpublished and often proprietary information. Review of specific tests, operators, and unpublished data are outside the scope of this evidence review and alternative sources exist. This evidence review focuses on the clinical validity and clinical utility.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Prospective Studies

A study by Tiersten et al (2009) used the Oncotech Extreme Drug Resistance (EDR) assay to examine whether chemotherapy resistance was an independent predictor of PFS in patients with stage III or IV ovarian cancer.¹⁹ Fifty-eight

eligible women were prospectively enrolled in this study; however, results from the EDR assay were not used to direct therapy. All women were treated with neoadjuvant chemotherapy and surgical cytoreduction followed by intraperitoneal chemotherapy. Evaluable EDR assay results were available for 22 patients. No difference in PFS was reported among patients who were defined as “resistant” or “nonresistant” to platinum or paclitaxel based on the EDR assay. Follow-up was not sufficient to measure OS. These data do not support the use of the EDR assay in predicting outcome and guiding patient management.

A review by Nagourney (2006) included 21 noncomparative studies using ex vivo programmed cell death assays.³ Selected studies correlated drug susceptibility findings for the ex vivo assay for those with an objective clinical response (complete or partial) and nonresponders (total 659 N=patients). Nagourney obtained aggregate positive values by the site of primary cancer: breast (82.9%), colon (80%), non-small-cell lung cancer (66.7%), gynecologic (77%), and small-cell lung cancer (50%). A 2012 study by this same investigator prospectively assessed 98 patients with non-small-cell lung cancer treated between 2003 and 2010.² Only 41 were eligible for inclusion and tested with the EVA/PCD assay to determine which chemotherapeutic drugs to use. Another 10 patients were excluded (5 due to insufficient cellular yield, 3 for resistance to all drugs tested, 2 due to physician’s choice), yielding only 31 patients who received the assay-recommended treatment. The authors compared results for these 31 patients treated using assay-directed chemotherapy with historical controls (not described) on the outcome of observed objective response rate (complete response and partial response). The objective response rate was 64.5% (95% confidence interval [CI], 46.9% to 78.9%) for the assay-directed chemotherapy group, which was significantly greater than the historic standard rate of 30% objective response ($p < 0.001$).

Retrospective Studies

Matsuo et al (2010) examined the relevance of extreme drug resistance in epithelial ovarian carcinomas.²⁰ Two hundred fifty-three records from the Oncotech database were identified for women with advanced stage ovarian cancer and from whom samples were collected at the time of the primary surgery. Tissue samples were cultured and tested for response to primary drugs (4 platinum- or taxane-based) and secondary drugs (eg, gemcitabine, topotecan, doxorubicin, etoposide, 5-fluorouracil). Paclitaxel showed the highest resistance rate. Other agents had resistance rates of less than 20%. Only 1 (0.4%) tumor showed complete resistance to all drugs tested, and 25% of tumors showed no resistance to any of the drugs. There was no statistical correlation between assay results and response to initial chemotherapy. Investigators acknowledged that the study, due to its retrospective and noncomparative design, was not sufficiently strong to validate the use of this assay in managing therapy. Potential confounding factors, described by investigators, included tumor heterogeneity and variations in resistance between primary tumor and metastases.

Another study by Matsuo et al (2009) evaluated the potential of the EDR assay to guide the selection of platinum- and taxane-based therapies for the management

of patients with advanced epithelial ovarian, fallopian, and peritoneal cancers.²¹ From the Oncotech database, 173 cases were identified. For all cases, tissue was collected at the time of cytoreductive therapy. The EDR assay was used to assess all samples, and tumors were classified as having low drug resistance, intermediate drug resistance, or extreme drug resistance. The 58 (33.5%) patients whose tumors had low drug resistance to both platinum and taxane showed statistically improved 5-year PFS and OS compared with the 115 (66.5%) patients who demonstrated intermediate or extreme drug resistance to platinum and/or taxane (5-year OS rates, 41.1% vs 30.9%, respectively; $p=0.014$). The 5-year OS rate in the 28 (16.2%) cases who had optimal cytoreduction with low drug resistance to both platinum and taxane (54.1%) was significantly better than that for the 62 (35.8%) cases who were suboptimally cyto-reduced with intermediate or extreme drug resistance to platinum and/or taxane (20.4%; $p<0.001$). Although the EDR assay was predictive for survival, assay results did not indicate a response to therapy with taxane or cisplatin. The investigators concluded that the EDR assay might be an independent predictor of PFS and OS; however, a prospective, randomized trial would be required to further assess its clinical utility in predicting response to taxane or platinum therapies.

In a smaller study (N=51) by Matsuo et al (2010), testing the predictive value of the EDR assay for uterine carcinosarcoma response to taxane and platinum, assessed 31 (60.8%) patients who received postoperative chemotherapy with at least a single agent and 17 (33.3%) received combination chemotherapy with platinum and taxane modalities.²² The overall response rate for the 17 combination chemotherapy cases was 70.6%. The presence of extreme drug resistance to platinum or taxane showed a significantly lower 1-year PFS rate (28.6% vs 100%, respectively; $p=0.01$) and lower 5-year OS rate (26.9% vs 57.1%, respectively; $p=0.033$). These data would indicate that use of an in vitro drug resistance assay might be predictive of response to chemotherapy response and survival outcome in advanced ovarian and uterine carcinosarcoma.

Matsuo et al (2010) also compared the rates of extreme drug resistance after cytoreductive therapy and neoadjuvant chemotherapy with the rates of extreme drug resistance after postoperative chemotherapy.²³ The goal of this study was not to test whether the EDR assay could direct therapeutic regimens. The findings suggested that platinum resistance was most common after neoadjuvant chemotherapy, while paclitaxel resistance was more prevalent after postoperative chemotherapy.

Karam et al (2009) retrospectively reviewed 377 patients with epithelial ovarian cancer to examine the effect of EDR assay-guided therapy on outcomes in the primary and recurrent setting.²⁴ The primary end points were time to progression, OS, and survival after recurrence. The patient population was heterogeneous, with a median age of 59 years (range, 24-89 years), and 30% of patients had completely resected tumors and were enrolled at varying tumor stages (Federation of Gynecologists and Obstetricians stages I, II, III, and IV in 7%, 4%, 78%, and 11%, respectively). Sixty-four percent of patients underwent a secondary cytoreductive surgery. Patients had an EDR assay sent either at the time of their

primary cytoreductive surgery (n=217) or at the time of disease recurrence (n=160). Predictors of survival included increasing age and greater volume of residual disease after cytoreductive surgery. EDR assay results analyzed for single agents or combinations of chemotherapies failed to predict patient outcomes independently whether the assay was performed at the time of the primary surgery or at recurrence.

Hetland et al (2012) studied primary platinum resistance in epithelial ovarian cancer patients with Federation of Gynecologists and Obstetricians stage III or IV disease.²⁵ Eighty-five biopsies from 58 patients were included. Resistance was assessed with a modified drug response assay including adenosine triphosphate (ATP)-based tumor chemosensitivity and EDR assay. Samples were tested for response to platinum, paclitaxel, and a combination of the drugs. Results from the assay were combined, and tumors were classified using a resistance index, which summarized the percentage of tumor growth inhibition for each drug concentration tested. All patients received a primary chemotherapy treatment of carboplatin, paclitaxel, or a combination of the 2 drugs. Platinum resistance, as defined by the risk index, was associated with significantly poorer PFS (median value, 3.9 months; 95% CI, 3.2 to 4.7 months) than platinum sensitivity (median PFS, 8.1 months; 95% CI, 3.7 to 12.4 months). Patients who had a partial response, stable disease or progressive disease were more resistant to platinum-based chemotherapy on the risk index score than those with a complete response (p=0.02). In a subgroup analysis of metastatic tumors, platinum resistance was not associated with PFS or clinical response. Response to paclitaxel or carboplatin plus paclitaxel was not associated with PFS or clinical response. In vitro response was not associated with OS in any group.

Section Summary: Clinically Valid

For chemoresistance assays, some retrospective and prospective correlational studies have suggested that these assays may be associated with chemotherapy response. However, prospective studies have not consistently demonstrated that chemoresistance assay results are associated with survival.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials (RCTs).

No studies comparing outcomes using assay-directed therapy with physician-chosen therapy were identified.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Because the clinical validity of chemosensitivity and chemoresistance assays has not been established, a chain of evidence cannot be constructed.

Section Summary: Chemoresistance Assays

Some retrospective and prospective studies have suggested that chemoresistance assays, particularly the EDR assay, may be associated with chemotherapy response. However, prospective studies have not consistently demonstrated that chemoresistance assay results are associated with survival. Furthermore, no comparative studies were identified that evaluated outcomes among patients managed with assay-directed therapy and those managed with physician-directed therapy. Large, randomized, prospective clinical studies comparing outcomes, including OS and disease-specific survival, QOL, and adverse events, between assay-directed therapy and physician-directed therapy, are needed.

Chemosensitivity Assays

Chemosensitivity assays are designed to select the most appropriate chemotherapy regimens for a given tumor type, and would therefore ideally be associated with high positive predictive values for clinical response. The critical type of evidence needed to establish the effectiveness of chemosensitivity assays would come from comparative studies of assay-guided therapy vs physician-directed therapy.

The TEC Assessment (2002)¹⁰ and systematic review by Samson et al (2004)¹¹ identified 9 comparative studies, 2 of which were randomized.²⁶⁻³⁴ Selected studies reported that significant advantages for assay-guided therapy regarding tumor response did not translate into survival differences. Response rate differences seen in other nonrandomized comparative studies may be attributable to bias or confounding, and survival outcomes were rarely reported.

Technically Reliable

Assessment of technical reliability focuses on specific tests and operators and requires review of unpublished and often proprietary information. Review of specific tests, operators, and unpublished data are outside the scope of this evidence review and alternative sources exist. This evidence review focuses on the clinical validity and clinical utility.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Prospective Studies

Kim et al (2010) reported on the results of a prospective, multicenter clinical trial designed to define the accuracy of the ATP-based chemotherapy response assay in

gastric cancer patients receiving paclitaxel and cisplatin chemotherapy, by comparing clinical response and the ATP assay results.³⁵ The primary study outcome was the accuracy of the ATP assay results, and the secondary outcome was finding the best method of defining in vitro chemosensitivity. Forty-eight patients with chemotherapy-naïve locally advanced or metastatic gastric cancer were treated with combination chemotherapy after a tissue specimen was obtained for the ATP assay. Tumor response was assessed by World Health Organization criteria using a computed tomography scan after every 2 cycles of chemotherapy. Laboratory technicians and physicians were blinded to the assay or clinical results. Thirty-six patients were evaluable for both in vitro and in vivo responses. Using a chemosensitivity index method, the specificity of the ATP assay was 95.7% (95% CI, 77.2% to 99.9%), sensitivity was 46.2% (95% CI, 19.2% to 74.9%), positive predictive value was 85.7% (95% CI, 42.1% to 99.6%), and NPV was 75.9% (95% CI, 55.1% to 89.3%). Median PFS was 4.2 months (95% CI, 3.4 to 5.0 months) and median OS was 11.8 months (95% CI, 9.7 to 13.8 months). The in vitro chemosensitive group showed a higher response rate (85.7%) than the chemoresistant group (24.1%; $p=0.005$). The authors concluded that the ATP assay could predict clinical response to paclitaxel and cisplatin chemotherapy with high accuracy in advanced gastric cancer.

Rutherford et al (2013) reported on results from a prospective, noninterventional, multicenter cohort study designed to assess whether the ChemoFX assay was predictive of outcomes among women ($N=335$) with histologically confirmed epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer.³⁶ Patients were treated with 1 of 15 study protocols; treating physicians were blinded to the ChemoFX assay result. Two hundred sixty-two (78.2% of total) patients had both available clinical follow-up data and a ChemoFX result. Cancer cells were classified based on the ChemoFX result as sensitive, intermediate, or resistant to each of several chemotherapeutic agents. Patients treated with an assay-sensitive regimen had a median PFS of 8.8 months compared with 5.9 months for those with assay-intermediate or -resistant regimens (hazard ratio [HR], 0.67; $p=0.009$). Mean OS was 37.5 months for patients treated with an assay-sensitive regimen and 23.9 months for those with assay-intermediate or -resistant regimens (HR=0.67, $p=0.010$).

In a follow-up analysis, Tian et al (2014) evaluated whether the ChemoFX assay could predict PFS by comparing the association when the assayed therapy matched the administered therapy (match) with the association when the assayed therapy was randomly selected (mismatch).³⁷ The authors generated a simulation in which the average prognostic value of assay results for multiple different therapies was generated using the assay results for a mismatch, in which the assay result for a treatment was randomly selected from the (up to) 15 designated therapies with equal probability for each patient. Based on 3000 repeated simulated resamplings, the mean HR for cases of mismatch was 0.81 (reported as 95% range, 0.66-0.99), which the authors suggested indicated that patients with a mismatch had less benefit when treated with an assay-sensitive therapy. Study strengths included the prospective design with physicians blinded to the assay results, which reduced the risk of bias in patient selection and outcomes

measurement. However, because the selection of chemotherapeutic agent was, by design, not influenced by the ChemoFX assay, the impact on health outcomes could not be determined.

Krivak et al (2014) reported on results of a subsequent prospective, observational, multicenter study to determine whether sensitivity to carboplatin and/or paclitaxel was associated with disease progression among patients with primary epithelial ovarian cancer following initial treatment with a platinum plus taxane regimen.³⁸ A total of 462 patients were enrolled, with 276 evaluable for analysis. Assay results for carboplatin and paclitaxel were available for 231 and 226 patients, respectively, with 44 (19.1%) patients identified as carboplatin-resistant and 49 (21.7%) identified as paclitaxel-resistant. Carboplatin-resistant patients were at a higher risk of disease progression than nonresistant patients (HR=1.87; 95% CI, 1.29 to 2.70; $p < 0.001$).

In a similar study design, Salom et al (2012) conducted a prospective, noninterventive, multicenter cohort study to assess whether the Microculture Kinetic (MiCK) assay (now called CorrectChemo) was predictive of outcomes among women with epithelial ovarian cancer.³⁹ Data from 150 women with any stage of cancer and specimens suitable for MiCK assay were included. Chemosensitivity was expressed as kinetic units following each dose of the drug in the MiCK assay and reported as mean, minimum, and maximum. For each patient, the "best" chemotherapy was defined as any single drug or combination of drugs in the patient's MiCK assay that had the highest kinetic units. Patients' regimens were at the discretion of treating physicians, who were blinded to the MiCK assay results. OS for stage III or IV disease was longer if patients received a chemotherapy considered "best" by the MiCK assay, compared with shorter OS for patients who received a chemotherapy that was not the best (HR=0.23, $p < 0.01$).

Jung et al (2013) conducted a single-center prospective study to determine whether sensitivity to paclitaxel and carboplatin, determined by using the Histoculture Drug Resistance Assay, was predictive of outcomes among 104 women with advanced epithelial ovarian cancer.⁴⁰ All patients had undergone initial surgery and were treated with paclitaxel and carboplatin therapy. Tumor cell sensitivity to the chemotherapy agents was classified as sensitive, intermediate, or resistant to paclitaxel, carboplatin, or both, based on the Histoculture Drug Resistance Assay. Patients whose tumors were sensitive to both drugs had a lower recurrence rate than those with resistance to both drugs (29.2% vs 69.8%, $p = 0.02$) and had a longer PFS (35 months vs 16 months; $p = 0.025$).

Zhang and Li (2015) evaluated ovarian epithelial cancer cells using an in vitro ATP tumor chemosensitivity assay.⁴¹ Specimens from 80 women with ovarian epithelial cancer who had undergone cytoreductive surgery were tested for sensitivity to 8 different treatments (paclitaxel, carboplatin, topotecan, gemcitabine, docetaxel, etoposide, bleomycin, 4-hydroperoxycyclophosphamide). Overall sensitivity, specificity, positive predictive value, and NPV were 88.6%, 77.8%, 83.0%, and 84.8%, respectively. Specimens from the lower stage (I-II) ovarian epithelial cancer had lower chemosensitivity than advanced stage (III). High to mildly

differentiated specimens had lower chemosensitivity than low differentiated specimens.

Retrospective Studies

A number of retrospective studies have evaluated the association between various chemosensitivity assays and clinical outcomes in several tumor types, most commonly epithelial ovarian cancer. Some representative studies are discussed next.

Tanigawa et al (2016) published a retrospective study evaluating the association between in vitro chemosensitivity results and relapse-free survival in 206 gastric cancer patients.⁴² The collagen gel droplet embedded culture drug sensitivity test is commercially available as a kit in Japan. All patients underwent surgery and were then treated with S-1 (tegafur/gimeracil/oteracil) chemotherapy. In vitro sensitivity of resected tumor specimens to fluorouracil was used as a surrogate of in vitro sensitivity to S-1 (this approach had been previously validated by the research group). Tumors were categorized as in vitro sensitive (responders) or in vitro insensitive (nonresponders). The median length of follow-up from the time of surgery was 3.2 years. Three-year relapse-free survival was significantly higher in the in vitro sensitive (responder) group (82.9%; 95% CI, 74.4% to 91.3%) than in the in vitro insensitive (nonresponder) group (63.4%; 95% CI, 54.7% to 72.1%; $p=0.001$).

Gallion et al (2006) retrospectively evaluated the association between ChemoFX test results and the treatment response for 256 patients with ovarian or peritoneal cancer treated with at least 1 cycle of postsurgical chemotherapy.⁴³ A subset of 135 patients had an exact match between drugs assayed and received; the rest had only a partial match. Predictive values were not reported or calculable. For the subset of 135, in a multivariable analysis, ChemoFX was an independent significant predictor ($p=0.006$) of PFS along with 2 other clinical variables. The HR for resistant vs sensitive was 2.9 (95% CI, 1.4 to 6.30) and was 1.7 (95% CI, 1.2 to 2.5) for resistant vs intermediate. The median PFS was 9 months for the resistant group, 14 months for the intermediate group, and had not been achieved at reporting for the sensitive group.

Herzog et al (2010)⁴⁴ included 147 patients from the Gallion study⁴³ and reported on 192 women with advanced stage primary ovarian cancer, 175 of whom had tumors tested for in vitro chemosensitivity to platinum therapy using ChemoFX. Tumors were classified as responsive, intermediately responsive, or nonresponsive to chemotherapy. Seventy-eight percent were categorized as responsive or intermediately responsive, and 22% were nonresponsive. Median OS was 72.5 months for patients with tumors categorized as responsive, 48.6 months for intermediately responsive, and 28.2 months for nonresponsive ($p=0.03$; HR=0.70; 95% CI, 0.50 to 0.97). The authors concluded that the results of chemosensitivity testing with a drug response marker for therapy were predictive of OS in patients with primary ovarian cancer.

In a smaller study, Grigsby et al (2013) retrospectively analyzed the association between pretreatment chemosensitivity to cisplatin and clinical outcomes in 33 women with cervical cancer.⁴⁵ Tumor cell sensitivity to cisplatin was categorized as responsive, intermediately responsive, or nonresponsive with the ChemoFX assay. Patients with responsive or intermediately responsive tumors had a 2-year recurrence-free survival of 87% compared with 58% for those with nonresponsive tumors ($p=0.036$).

Lee et al (2012) conducted a retrospective study of Histoculture Drug Resistance Assay in 79 patients with ovarian cancer.⁴⁶ Tissue samples were assessed for 11 chemotherapeutic agents and found the highest inhibition rates in carboplatin (49.2%), topotecan (44.7%), and belotecan (39.7%). These inhibition rates were higher than with cisplatin (34.7%), the traditional drug used to treat epithelial ovarian cancer. Outcomes for a subset of 37 patients with Federation of Gynecologists and Obstetricians stages II through IV epithelial ovarian serous adenocarcinoma who had been treated with at least 3 cycles of carboplatin chemotherapy were assessed with carboplatin-sensitive and -resistant patients. Multiple comparisons and regression analyses established a 40% inhibition rate cutoff value in response to 50 $\mu\text{g/mL}$ carboplatin to determine sensitivity or resistance. This selected cutoff had a disease-free survival of 23.2 months (95% CI, 6.3 to 55.3 months) and 13.8 months (95% CI, 4.9 to 35.6 months) in the carboplatin-sensitive and carboplatin-resistant groups, respectively ($p<0.05$). Mean OS between groups did not differ significantly (carboplatin-sensitive patients, 60.4 months vs carboplatin-resistant patients, 37.3 months; $p=0.621$).

Strickland et al (2013) retrospectively evaluated the association between chemosensitivity and anthracyclines, measured by the drug-induced apoptosis MiCK assay, among 109 patients with adult-onset acute myelogenous leukemia.⁴⁷ Patients were treated with a "7 plus 3" chemotherapy regimen. Chemosensitivity was expressed as maximal kinetic units following each dose of the drug in the MiCK assay. Receiver operator characteristic curve analysis and logistic regression were used to determine the optimal cutoff for chemosensitivity response to discriminate between chemoresponder and nonresponder. Patients determined to be chemoresponders to idarubicin were more likely to have a complete response to chemotherapy (72%) than those who were nonresponders ($p=0.01$). Data for the patient cohort were collected over 14 years, from 1996 to 2010, which would limit the generalizability of the results to current chemotherapy regimens.

Other retrospective studies have evaluated chemosensitivity results as measured by other assay types. Von Heideman et al (2014) evaluated the semi-automated fluorometric microculture cytotoxicity assay in 112 patients (125 samples) with ovarian cancer and concluded that samples from patients with clinical response were more sensitive to most drugs than samples from nonresponding patients.⁴⁸

Section Summary: Clinically Valid

For chemosensitivity assays, the evidence includes retrospective and prospective correlational studies. These studies of several different chemosensitivity assays

have suggested that patients whose tumors have higher chemosensitivity have better outcomes.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

A small number of nonrandomized studies have evaluated differences in outcomes between patients treated with assay-directed therapy and patient given with physician-chosen therapy.

Bosserman et al (2015) published a prospective nonblinded study to determine if physicians who use the results from MiCK assays on breast cancer specimens have better patient outcomes than physicians who do not.⁴⁹ Tumor samples were extracted from 30 women with recurrent or metastatic breast cancer and submitted for the MiCK drug-induced apoptosis assay. Results were available to physicians within 72 hours after the biopsy. Physicians could use or not use the test results to determine therapy. Most physicians (22/30) used the assay results to select the chemotherapy regimens for their patients. Of those using the assay results, 15 physicians changed the original treatment plans for their patients. Among physicians who did not use the assay results, reasons given included: patient refused the most active drugs indicated by the assay (4 patients), the physician did not want to use most active drugs indicated by the assay (2 patients), and unstated (2 patients). Complete response, partial response, and disease control were more frequently experienced in patients whose physicians used the assay results compared with patients whose physicians did not use the assay results ($p=0.04$). Time to recurrence was significantly longer in patients whose physicians used the assay (7.4 months) compared with patients whose physicians did not (2.2 months). OS did not differ significantly between patients whose physicians used the assay (16.8 months) and patients whose physicians did not (13.1 months).

Ugurel et al (2006) conducted a nonrandomized, prospective phase 2 trial, of 53 evaluable patients with metastatic melanoma.⁵⁰ Biopsy samples from each patient were sent to a laboratory for chemosensitivity testing using the ATP tumor chemosensitivity assay. Patients then received assay-directed therapy with the drug or drug combination that had the highest in vitro sensitivity. Median follow-up was 19 months. The study found a 36% complete and partial response rate in patients with chemosensitive tumors compared with 16% in those with chemoresistant tumors.

In a case-control retrospective trial, Moon et al (2009) retrospectively compared ATP assay-based guided chemotherapy with empirical chemotherapy in unresectable non-small-cell lung cancer.⁵¹ All patients who received ATP assay-guided platinum-based doublet chemotherapy as first-line therapy received platinum-based chemotherapy combined with a nonplatinum drug, regardless of their in vitro platinum sensitivity; 14 patients had platinum-sensitive disease, and 13 were platinum-resistant. Ninety-three matched controls (matched for performance status, stage, and chemotherapy regimen) were selected from a retrospective review of a database. In the empirical group, a nonplatinum drug was chosen, depending on physicians' discretion, along with a platinum agent determined by renal function and performance status. The primary end point was clinical response rate, assessed every 2 cycles of chemotherapy using the Response Evaluation Criteria in Solid Tumors criteria. The secondary end points were PFS and OS. Response and survival rates in both groups did not differ statistically. By ATP assay, the platinum-sensitive subgroup (71%) showed a higher response rate than the empirical group (38%; $p=0.02$), but PFS and OS differences between groups were not statistically significant.

In a nonrandomized prospective study (N=64), Iwahashi et al (2005) reported on outcomes of chemosensitivity-guided chemotherapy compared with standard chemotherapy and no chemotherapy in patients with advanced gastric cancer who had undergone a gastrectomy.⁵² Among patients with stage IV gastric cancer, the 5-year OS rate was 38% in the chemosensitivity-guided chemotherapy group and 0% in the standard chemotherapy and no chemotherapy groups. Among patients with para-aortic node involvement, survival was significantly greater in the chemosensitivity-guided group than in with the standard and no chemotherapy groups. However, survival was equivalent between the groups when there was no para-aortic node involvement.

Cree et al (2007) reported on a prospective RCT of chemosensitivity assay-directed chemotherapy vs physician's choice in patients with recurrent platinum-resistant ovarian cancer.⁵³ The primary aim of this RCT was to determine response and PFS rates following chemotherapy in patients treated based on an ATP-based tumor chemosensitivity assay or physician-choice. A total of 180 patients were randomized to assay-directed therapy (n=94) or to physician-choice chemotherapy (n=86). Median follow-up at analysis was 18 months; the response was assessable in 147 (82%) patients: 31.5% achieved a partial or complete response in the physician-choice group compared with 40.5% in the assay-directed group (26% vs 31% by intention-to-treat analysis, respectively). Intention-to-treat analysis showed a median PFS of 93 days in the physician-choice group and 104 days in the assay-directed group (HR=0.8; $p=0.14$). No difference was reported in OS between groups, although 12 (41%) of 39 patients who crossed over from the physician-choice arm obtained a response. Increased use of combination therapy was seen in the physician-choice arm during the trial as a result of the observed effects of assay-directed therapy in patients. The authors concluded that this small RCT suggested a trend toward improved response and PFS for assay-directed treatment and that chemosensitivity testing might provide useful information in some patients with ovarian cancer. They also

noted that the ATP-based tumor chemosensitivity assay would remain an investigational method in this condition.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Because the clinical validity of chemosensitivity and chemoresistance assays has not been established, a chain of evidence cannot be constructed.

Section Summary: Chemosensitivity Assays

The most direct evidence related to the effectiveness of chemosensitivity assays in the management of patients with cancer comes from several studies that compare outcomes for patients managed using an ATP-based tumor chemosensitivity assay with those managed using standard care, including an RCT. Although some improvements in tumor response were noted, no differences between OS or PFS were seen. A number of retrospective and prospective studies of several different chemosensitivity assays, including the ATP-based tumor chemosensitivity assay, the CorrectChemo assay, and the ChemoFX assay, have suggested that patients whose tumors have higher chemosensitivity have better outcomes. However, additional studies to determine whether the clinical use of in vitro chemosensitivity testing leads to better outcomes are needed.

Summary of Evidence

For individuals who have cancer who are initiating chemotherapy who receive chemoresistance assays, the evidence includes correlational observational studies. Relevant outcomes are overall survival, disease-specific survival, test accuracy and validity, and quality of life. Some retrospective and prospective correlational studies have suggested that chemoresistance assays may be associated with chemotherapy response. However, prospective studies have not consistently demonstrated that chemoresistance assay results are associated with survival. Furthermore, no studies were identified that compared outcomes for patients managed using assay-directed therapy with those managed using physician-directed therapy. Large, randomized, prospective clinical studies comparing overall survival are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have cancer who are initiating chemotherapy who receive chemosensitivity assays, the evidence includes an RCT, nonrandomized studies, and correlational observational studies. Relevant outcomes are overall survival, disease-specific survival, test accuracy and validity, and quality of life. The most direct evidence on the effectiveness of chemosensitivity assays in the management of patients with cancer comes from several studies comparing outcomes for patients managed using a chemosensitivity assay with those managed using standard care, including an RCT. Although some improvements in tumor response were noted in the randomized trial, there were no differences in survival outcomes. One small nonrandomized study reported improved overall

survival in patients receiving chemosensitivity-guided therapy compared with patients receiving standard chemotherapy. A number of retrospective and prospective studies of several different chemosensitivity assays have suggested that patients whose tumors have higher chemosensitivity have better outcomes. Currently, additional studies to determine whether the clinical use of in vitro chemosensitivity testing leads to improvements in overall survival are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

Supplemental Information

Practice Guidelines and Position Statements

National Comprehensive Cancer Network

Current National Comprehensive Cancer Network (NCCN) guidelines for the treatment of epithelial ovarian cancer, fallopian tube cancer, and primary peritoneal cancer (v.2.2018) state that “Chemosensitivity/resistance and/or other biomarker assays are being used in some NCCN Member Institutions for decisions related to future chemotherapy in situations where there are multiple equivalent chemotherapy options available. The current level of evidence is not sufficient to supplant standard-of-care chemotherapy (category 3).”⁵⁴

NCCN guidelines for the treatment of gastric cancer (v.2.2018)⁵⁵ and uterine neoplasms (v.2.2018)⁵⁶ do not discuss the use of chemoresistance or chemosensitivity assays as part of cancer management.

American Society of Clinical Oncology

The updated 2011 American Society of Clinical Oncology clinical guidelines on the use of chemotherapy sensitivity and resistance assays did not recommend the use of chemotherapy sensitivity and resistance assays unless in a clinical trial setting.⁵⁷

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

An ongoing trials that might influence this review is listed in Table 1.

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing NCT02580253	Adjuvant Chemotherapy Based on the Adenosine	300	Dec 2018

Triphosphate Tumor Chemosensitivity Assay for
Hepatocellular Carcinoma After Liver
Transplantation

NCT03133273^a	Study of the Therapeutic Response and survival of Patients with Metastatic Colorectal Cancer (Stage IV) and Treated According to the Guidelines of a Chemosensitivity Test, Oncogramme®	230	Jul 2020
--------------------------------	---	-----	----------

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

References

- Bird MC, Godwin VA, Antrobus JH, et al. Comparison of in vitro drug sensitivity by the differential staining cytotoxicity (DiSC) and colony-forming assays. *Br J Cancer*. Apr 1987;55(4):429-431. PMID 3580265
- Nagourney RA, Blitzer JB, Shuman RL, et al. Functional profiling to select chemotherapy in untreated, advanced or metastatic non-small cell lung cancer. *Anticancer Res*. Oct 2012;32(10):4453-4460. PMID 23060572
- Nagourney RA. Ex vivo programmed cell death and the prediction of response to chemotherapy. *Curr Treat Options Oncol*. Mar 2006;7(2):103-110. PMID 16455021
- Csoka K, Larsson R, Tholander B, et al. Cytotoxic drug sensitivity testing of tumor cells from patients with ovarian carcinoma using the fluorometric microculture cytotoxicity assay (FMCA). *Gynecol Oncol*. Aug 1994;54(2):163-170. PMID 7520407
- Yung WK. In vitro chemosensitivity testing and its clinical application in human gliomas. *Neurosurg Rev*. Jan 1989;12(3):197-203. PMID 2682352
- Kern DH, Weisenthal LM. Highly specific prediction of antineoplastic drug resistance with an in vitro assay using suprapharmacologic drug exposures. *J Natl Cancer Inst*. Apr 4 1990;82(7):582-588. PMID 2313735
- Anticancer Inc. Histoculture Drug Response Assay - HDRA. n.d.; http://www.anticancer.com/HDRA_ref.html. Accessed June 5, 2018.
- Helomics. ChemoFx Chemoresponse Marker. n.d.; <https://www.helomics.com/chemoresponse-patients>. Accessed June 5, 2018.
- Brower SL, Fensterer JE, Bush JE. The ChemoFx assay: an ex vivo chemosensitivity and resistance assay for predicting patient response to cancer chemotherapy. *Methods Mol Biol*. 2008;414:57-78. PMID 18175812
- Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Chemotherapy Sensitivity and Resistance Assays. TEC Assessments. 2002;17(12). PMID 12166470
- Samson DJ, Seidenfeld J, Ziegler K, et al. Chemotherapy sensitivity and resistance assays: a systematic review. *J Clin Oncol*. Sep 1 2004;22(17):3618-3630. PMID 15289487
- Brown E, Markman M. Tumor chemosensitivity and chemoresistance assays. *Cancer*. Mar 15 1996;77(6):1020-1025. PMID 8635118
- Eltabbakh GH, Piver MS, Hempling RE, et al. Correlation between extreme drug resistance assay and response to primary paclitaxel and cisplatin in patients with epithelial ovarian cancer. *Gynecol Oncol*. Sep 1998;70(3):392-397. PMID 9790793
- Eltabbakh GH. Extreme drug resistance assay and response to chemotherapy in patients with primary peritoneal carcinoma. *J Surg Oncol*. Mar 2000;73(3):148-152. PMID 10738268
- Mehta RS, Bornstein R, Yu IR, et al. Breast cancer survival and in vitro tumor response in the extreme drug resistance assay. *Breast Cancer Res Treat*. Apr 2001;66(3):225-237. PMID 11510694
- Holloway RW, Mehta RS, Finkler NJ, et al. Association between in vitro platinum resistance in the EDR assay and clinical outcomes for ovarian cancer patients. *Gynecol Oncol*. Oct 2002;87(1):8-16. PMID 12468336
- Ellis RJ, Fabian CJ, Kimler BF, et al. Factors associated with success of the extreme drug resistance assay in primary breast cancer specimens. *Breast Cancer Res Treat*. Jan 2002;71(2):95-102. PMID 11881914
- Loizzi V, Chan JK, Osann K, et al. Survival outcomes in patients with recurrent ovarian cancer who were treated with chemoresistance assay-guided chemotherapy. *Am J Obstet Gynecol*. Nov 2003;189(5):1301-1307. PMID 14634558

19. Tiersten AD, Moon J, Smith HO, et al. Chemotherapy resistance as a predictor of progression-free survival in ovarian cancer patients treated with neoadjuvant chemotherapy and surgical cytoreduction followed by intraperitoneal chemotherapy: a Southwest Oncology Group Study. *Oncology*. Feb 2009;77(6):395-399. PMID 20130422
20. Matsuo K, Eno ML, Im DD, et al. Clinical relevance of extent of extreme drug resistance in epithelial ovarian carcinoma. *Gynecol Oncol*. Jan 2010;116(1):61-65. PMID 19840886
21. Matsuo K, Bond VK, Eno ML, et al. Low drug resistance to both platinum and taxane chemotherapy on an in vitro drug resistance assay predicts improved survival in patients with advanced epithelial ovarian, fallopian and peritoneal cancer. *Int J Cancer*. Dec 1 2009;125(11):2721-2727. PMID 19530239
22. Matsuo K, Bond VK, Im DD, et al. Prediction of chemotherapy response with platinum and taxane in the advanced stage of ovarian and uterine carcinosarcoma: a clinical implication of in vitro drug resistance assay. *Am J Clin Oncol*. Aug 2010;33(4):358-363. PMID 19875949
23. Matsuo K, Eno ML, Im DD, et al. Chemotherapy time interval and development of platinum and taxane resistance in ovarian, fallopian, and peritoneal carcinomas. *Arch Gynecol Obstet*. Feb 2010;281(2):325-328. PMID 19455347
24. Karam AK, Chiang JW, Fung E, et al. Extreme drug resistance assay results do not influence survival in women with epithelial ovarian cancer. *Gynecol Oncol*. Aug 2009;114(2):246-252. PMID 19500821
25. Hetland TE, Kaern J, Skrede M, et al. Predicting platinum resistance in primary advanced ovarian cancer patients with an in vitro resistance index. *Cancer Chemother Pharmacol*. May 2012;69(5):1307-1314. PMID 22302409
26. Cortazar P, Gazdar AF, Woods E, et al. Survival of patients with limited-stage small cell lung cancer treated with individualized chemotherapy selected by in vitro drug sensitivity testing. *Clin Cancer Res*. May 1997;3(5):741-747. PMID 9815744
27. Gazdar AF, Steinberg SM, Russell EK, et al. Correlation of in vitro drug-sensitivity testing results with response to chemotherapy and survival in extensive-stage small cell lung cancer: a prospective clinical trial. *J Natl Cancer Inst*. Jan 17 1990;82(2):117-124. PMID 2152944
28. Kurbacher CM, Cree IA, Bruckner HW, et al. Use of an ex vivo ATP luminescence assay to direct chemotherapy for recurrent ovarian cancer. *Anticancer Drugs*. Jan 1998;9(1):51-57. PMID 9491792
29. Shaw GL, Gazdar AF, Phelps R, et al. Individualized chemotherapy for patients with non-small cell lung cancer determined by prospective identification of neuroendocrine markers and in vitro drug sensitivity testing. *Cancer Res*. Nov 1 1993;53(21):5181-5187. PMID 8221655
30. Shaw GL, Gazdar AF, Phelps R, et al. Correlation of in vitro drug sensitivity testing results with response to chemotherapy and survival: comparison of non-small cell lung cancer and small cell lung cancer. *J Cell Biochem Suppl*. Jan 1996;24:173-185. PMID 8806100
31. Von Hoff DD, Kronmal R, Salmon SE, et al. A Southwest Oncology Group study on the use of a human tumor cloning assay for predicting response in patients with ovarian cancer. *Cancer*. Jan 1 1991;67(1):20-27. PMID 1985717
32. Von Hoff DD, Sandbach JF, Clark GM, et al. Selection of cancer chemotherapy for a patient by an in vitro assay versus a clinician. *J Natl Cancer Inst*. Jan 17 1990;82(2):110-116. PMID 2403593
33. Wilbur DW, Camacho ES, Hilliard DA, et al. Chemotherapy of non-small cell lung carcinoma guided by an in vitro drug resistance assay measuring total tumour cell kill. *Br J Cancer*. Jan 1992;65(1):27-32. PMID 1310250
34. Xu JM, Song ST, Tang ZM, et al. Predictive chemotherapy of advanced breast cancer directed by MTT assay in vitro. *Breast Cancer Res Treat*. Jan 1999;53(1):77-85. PMID 10206075
35. Kim JH, Lee KW, Kim YH, et al. Individualized tumor response testing for prediction of response to Paclitaxel and Cisplatin chemotherapy in patients with advanced gastric cancer. *J Korean Med Sci*. May 2010;25(5):684-690. PMID 20436702
36. Rutherford T, Orr J, Jr., Grendys E, Jr., et al. A prospective study evaluating the clinical relevance of a chemoresponse assay for treatment of patients with persistent or recurrent ovarian cancer. *Gynecol Oncol*. Nov 2013;131(2):362-367. PMID 23954900
37. Tian C, Sargent DJ, Krivak TC, et al. Evaluation of a chemoresponse assay as a predictive marker in the treatment of recurrent ovarian cancer: further analysis of a prospective study. *Br J Cancer*. Aug 26 2014;111(5):843-850. PMID 25003664

38. Krivak TC, Lele S, Richard S, et al. A chemoresponse assay for prediction of platinum resistance in primary ovarian cancer. *Am J Obstet Gynecol*. Jul 2014;211(1):68 e61-68. PMID 24530815
39. Salom E, Penalver M, Homesley H, et al. Correlation of pretreatment drug induced apoptosis in ovarian cancer cells with patient survival and clinical response. *J Transl Med*. Aug 08 2012;10:162. PMID 22873358
40. Jung PS, Kim DY, Kim MB, et al. Progression-free survival is accurately predicted in patients treated with chemotherapy for epithelial ovarian cancer by the histoculture drug response assay in a prospective correlative clinical trial at a single institution. *Anticancer Res*. Mar 2013;33(3):1029-1034. PMID 23482777
41. Zhang J, Li H. Heterogeneity of tumor chemosensitivity in ovarian epithelial cancer revealed using the adenosine triphosphate-tumor chemosensitivity assay. *Oncol Lett*. May 2015;9(5):2374-2380. PMID 26137074
42. Tanigawa N, Yamaue H, Ohyama S, et al. Exploratory phase II trial in a multicenter setting to evaluate the clinical value of a chemosensitivity test in patients with gastric cancer (JACCRO-GC 04, Kubota memorial trial). *Gastric Cancer*. Apr 2016;19(2):350-360. PMID 26385385
43. Gallion H, Christopherson WA, Coleman RL, et al. Progression-free interval in ovarian cancer and predictive value of an ex vivo chemoresponse assay. *Int J Gynecol Cancer*. Jan-Feb 2006;16(1):194-201. PMID 16445633
44. Herzog TJ, Krivak TC, Fader AN, et al. Chemosensitivity testing with ChemoFx and overall survival in primary ovarian cancer. *Am J Obstet Gynecol*. Jul 2010;203(1):68 e61-66. PMID 20227055
45. Grigsby PW, Zigelboim I, Powell MA, et al. In vitro chemoresponse to cisplatin and outcomes in cervical cancer. *Gynecol Oncol*. Jul 2013;130(1):188-191. PMID 23583416
46. Lee JH, Um JW, Lee JH, et al. Can immunohistochemistry of multidrug-resistant proteins replace the histoculture drug response assay in colorectal adenocarcinomas? *Hepatogastroenterology*. Jun 2012;59(116):1075-1078. PMID 22580657
47. Strickland SA, Raptis A, Hallquist A, et al. Correlation of the microculture-kinetic drug-induced apoptosis assay with patient outcomes in initial treatment of adult acute myelocytic leukemia. *Leuk Lymphoma*. Mar 2013;54(3):528-534. PMID 22924433
48. von Heideman A, Tholander B, Grundmark B, et al. Chemotherapeutic drug sensitivity of primary cultures of epithelial ovarian cancer cells from patients in relation to tumour characteristics and therapeutic outcome. *Acta Oncol*. Feb 2014;53(2):242-250. PMID 23713890
49. Bosserman L, Rogers K, Willis C, et al. Application of a drug-induced apoptosis assay to identify treatment strategies in recurrent or metastatic breast cancer. *PLoS One*. May 29 2015;10(5):e0122609. PMID 26024531
50. Ugurel S, Schadendorf D, Pfohler C, et al. In vitro drug sensitivity predicts response and survival after individualized sensitivity-directed chemotherapy in metastatic melanoma: a multicenter phase II trial of the Dermatologic Cooperative Oncology Group. *Clin Cancer Res*. Sep 15 2006;12(18):5454-5463. PMID 17000680
51. Moon YW, Sohn JH, Kim YT, et al. Adenosine triphosphate-based chemotherapy response assay (ATP-CRA)-guided versus empirical chemotherapy in unresectable non-small cell lung cancer. *Anticancer Res*. Oct 2009;29(10):4243-4249. PMID 19846981
52. Iwahashi M, Nakamori M, Nakamura M, et al. Individualized adjuvant chemotherapy guided by chemosensitivity test sequential to extended surgery for advanced gastric cancer. *Anticancer Res*. Sep-Oct 2005;25(5):3453-3459. PMID 16101163
53. Cree IA, Kurbacher CM, Lamont A, et al. A prospective randomized controlled trial of tumour chemosensitivity assay directed chemotherapy versus physician's choice in patients with recurrent platinum-resistant ovarian cancer. *Anticancer Drugs*. Oct 2007;18(9):1093-1101. PMID 17704660
54. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Ovarian Cancer Including Fallopian Tube Cancer and Primary Peritoneal Cancer. Version 2.2018. https://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf. Accessed June 7, 2018.
55. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Gastric Cancer. Version 2.2018. https://www.nccn.org/professionals/physician_gls/pdf/gastric.pdf. Accessed June 7, 2018.

56. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Uterine Neoplasms. Version 2.2018. https://www.nccn.org/professionals/physician_gls/pdf/uterine.pdf. Accessed June 7, 2018.
57. Burstein HJ, Mangu PB, Somerfield MR, et al. American Society of Clinical Oncology clinical practice guideline update on the use of chemotherapy sensitivity and resistance assays. J Clin Oncol. Aug 20 2011;29(24):3328-3330. PMID 21788567

Billing Coding/Physician Documentation Information

- 81535** Oncology (gynecologic), live tumor cell culture and chemotherapeutic response by DAPI stain and morphology, predictive algorithm reported as a drug response score; first single drug or drug combination
- 81536** Oncology (gynecologic), live tumor cell culture and chemotherapeutic response by DAPI stain and morphology, predictive algorithm reported as a drug response score; each additional single drug or drug combination (List separately in addition to code for primary procedure)
- 88358** Morphometric analysis; tumor (eg, DNA ploidy)
- 88305** Level IV Surgical pathology, gross and microscopic examination Abortion
- 88104** Cytopathology, fluids, washings or brushings, except cervical or vaginal; smears with interpretation
- 87230** Toxin or antitoxin assay, tissue culture (eg, Clostridium difficile toxin)
- 88313** Special stains (List separately in addition to code for primary service); Group II, all other, (eg, iron, trichrome), except immunocytochemistry and immunoperoxidase stains, each
- 89050** Cell count, miscellaneous body fluids (eg, cerebrospinal fluid, joint fluid), except blood;

ICD-10 Codes

- C00.0-** Malignant neoplasms code range
- C96.9**

The extreme drug resistance assay is a multistep laboratory procedure identified by the CPT codes listed above.

Effective January 1, 2016, there will be specific CPT codes for ChemoFX®: 81535 Oncology (gynecologic), live tumor cell culture and chemotherapeutic response by DAPI stand and morphology, predictive algorithm reported as a drug response score; first single drug or drug combination 81536 each additional single drug or drug combination (List separately in addition to code for primary procedure).

Additional Policy Key Words

N/A

Policy Implementation/Update Information

- 10/1/88 New policy.
- 10/1/00 No policy statement changes.
- 10/1/01 No policy statement changes.
- 10/1/02 No policy statement changes.

- 10/1/03 No policy statement changes.
 - 10/1/04 No policy statement changes.
 - 10/1/05 No policy statement changes.
 - 10/1/06 No policy statement changes.
 - 10/1/07 No policy statement changes.
 - 10/1/08 No policy statement changes.
 - 10/1/09 No policy statement changes.
 - 10/1/10 No policy statement changes.
 - 10/1/11 No policy statement changes.
 - 10/1/12 No policy statement changes.
 - 10/1/13 No policy statement changes.
 - 10/1/14 No policy statement changes.
 - 10/1/15 "ChemoFx" and "CorrectChemo" added to the list of investigational chemosensitivity assays; policy statements otherwise unchanged.
 - 10/1/16 No policy statement changes.
 - 10/1/17 No policy statement changes.
 - 10/1/18 CorrectChemo assay removed from the second policy statement; intent of statements unchanged.
-

State and Federal mandates and health plan contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. The medical policies contained herein are for informational purposes. The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents Blue KC and are solely responsible for diagnosis, treatment and medical advice. No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, photocopying, or otherwise, without permission from Blue KC.